

| | PCR | isolation | antigen detection | | serology | total this 2/52 | total prev 2/52 | comments |
|-----------------------------------|-----|-----------|-------------------|-------|----------|--------------------|--------------------|--|
| | | | IF | other | | | | |
| Human metapneumovirus | 5 | | | | | 5 | 3 | |
| Influenza A | 8 | | | | 10 | 18 | 18 | |
| Influenza B | 2 | | | | 3 | 5 | 3 | |
| Legionella spp | | | | 3* | 18 | 21 | 5 | * Urinary Ag |
| M. tuberculosis | | 10 | | | | 10 | 14 | |
| M. bovis BCG | | | | | | | | |
| Other significant mycobacteria | | 3* | | | | 3 | | * M. abscessus, M. marinum (x2) |
| Mycoplasma | | | | | 7 | 7 | 6 | |
| Parainfluenza | 7 | | | | | 7 | 3 | |
| Pertussis | | | | | | | 3 | |
| Picornavirus | 17 | | | | | 17 | 20 | |
| Psittacosis | | | | | 5 | 5 | 5 | |
| Q fever | | | | | | | 1 | |
| RSV | 2 | | | | | 2 | | |
| Skin/Mucosal Lesions | | | | | | | | |
| Herpes simplex | 13 | | | | | 13 | 8 | |
| M. ulcerans | 2 | | | | | 2 | 4 | |
| Poxvirus (M. contagiosum-like) | | | | | | | | |
| Uro-genital Pathogens | | | | | | | | |
| Chlamydia | 33 | | | | | 33 | 31 | |
| Herpes simplex | 25 | | | | | 25 | 22 | |
| N. gonorrhoeae | 7 | 7 | | | | 14 | 26 | 5x urogenital, 2x rectal, 6x throat, 1x urine |
| Syphilis | 8 | | | | 32* | 40 | 36 | *8 RPR > 16 |
| Varicella zoster | 1 | | | | | 1 | 2 | |