

## Report No. 1: Week ending 6 May 2018


### Overview: The 2018 influenza season so far

Influenza surveillance indicates activity is at **baseline levels**.

The number of notified **laboratory confirmed influenza** cases for the year to 6 May is **higher** than at the same time in 2017. Cases notified year-to-date have been type A (47%) and type B (49%).

Seven VicSPIN swabs were received this week, one positive for type A (H1N1) and one positive for influenza B. VicSPIN ILI rates are at baseline levels.

*Figure 1. Routinely notified influenza cases and VicSPIN and National Home Doctor Service (NHDS) ILI consultation proportions, Victoria, 2007-2018*



### National Home Doctor Service (Victoria)

- The ILI proportion for the week was 4.4 ILI cases per 1,000 patients seen (Figure 1).
- ILI proportions are at baseline levels.

## Victorian Sentinel Practice Influenza Network

- The total influenza-like illness (ILI) proportion measured by VicSPIN is at baseline levels, at 0.7 ILI cases per 1,000 patients seen this week. (Figures 1 & 2).
- Seven swabs were received this week, one positive for type A (H1N1) and one positive for influenza B.

*Table 1. Summary of reporting from general practice sentinel sites, week ending 6 May 2018*

	Total
Number (%) of general practitioners notifying	75 (78%)
Total patients seen	5,754
Total patients with ILI	4
Consultation rate for ILI (per 1,000 patients)	0.7
Number of specimens received	7
Number (%) positive for influenza	2 (25%)

*Figure 2. VicSPIN ILI consultation proportions, Victoria, 2018*





Figure 3. VicSPIN respiratory virus detections and percent influenza positive, Victoria, 2018



### Observations from other influenza surveillance systems

#### Notified cases of laboratory confirmed influenza

- A total of 1,899 cases of laboratory confirmed influenza have been processed so far in 2018 (Figure 1), compared with 1,385 in 2017, although this may be revised up as data are updated.
- Cases notified year-to-date have been type A (47%) and type B (49%). Type A cases had a median age of 48 years and type B cases median age 50 years.

#### World Health Organization Collaborating Centre for Reference and Research on Influenza

- The strains included in the 2018 quadrivalent influenza vaccine are: A/Michigan/45/2015 (H1N1)pdm09-like virus; A/Singapore/INFIMH-16-0019/2016 (H3N2)-like virus; B/Phuket/3073/2013-like virus with the quadrivalent vaccine including B/Brisbane/60/2008-like virus as well.
- No Victorian isolates have been strain-typed by the WHO so far in 2018.
- No isolate has shown resistance to Zanamivir, among the 80 Australian isolates tested so far in 2018.