

Background

The noroviruses are the most common cause of outbreaks of non-bacterial gastroenteritis as well as being an important cause of sporadic gastroenteritis. Although norovirus-associated gastroenteritis was once referred to as “winter vomiting disease”, it is now known that norovirus-associated gastroenteritis occurs throughout the year. Furthermore, a recent study of norovirus outbreaks in Victoria over the period 2001-2010 showed that there had been a dramatic increase in norovirus outbreaks in the period 2006-2010 compared to 2001-2005.¹

The available evidence indicates this increase was not a function of enhanced surveillance procedures but most probably related to the emergence of more virulent strains of norovirus and also to environmental conditions which facilitated the spread of the virus. This report, based on laboratory detections of norovirus and general practitioner gastroenteritis presentations, is intended to provide an indication of norovirus and gastroenteritis activity in Victoria.

Summary

Routine testing at the Victorian Infectious Diseases Reference Laboratory (VIDRL) includes detections of norovirus from outbreaks and sporadic cases. Because testing tends to focus on gastroenteritis outbreaks in aged-care facilities, detection of norovirus may be biased towards older age groups. The number of norovirus detections has increased this month, with the majority of cases in the 50-79 and 80+ age groups (Figure 1). It is too soon to say whether this increase may be the start of the annual peak in cases, or represents seasonal variation.

The Melbourne Medical Deputising Service (MMDS), an out-of-hours general practitioner locum service, provides data on gastroenteritis-like illness consultations. By the end of June the rate of gastroenteritis per 1000 consultations had risen to 58.8 (Figure 2). As seen previously rates were highest in the 18-<65 year old age group.

These data suggest that gastroenteritis activity fell from a peak earlier in the year and may again be rising in Victoria. The increase in activity observed in late 2012/ early 2013 may be due to the emergence of the genotype II.4 2012 Sydney variant. Increases in norovirus activity due to this genotype have been recorded in the United Kingdom, France, Denmark and New Zealand^{2, 3, 4}.

¹LD Bruggink, K Sturge, J Gaston, J Gregory, MG Catton, JA Marshall *Patterns of norovirus-associated gastroenteritis outbreaks in Victoria 2001–2010* Victorian Infectious Diseases Bulletin Volume 14 Issue 3 September 2011

²van Beek J, Ambert-Balay K, Botteldoorn N, Eden JS, Fonager J, Hewitt J, Iritani N, Kroneman A, Vennema H, Vinjé J, White PA, Koopmans M, on behalf of NoroNet. Indications for worldwide increased norovirus activity associated with emergence of a new variant of genotype II.4, late 2012. *Euro Surveill.* 2013;18(1):pii=20345. Available online: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=20345>

³Fonager J, Hindbæk LS, Fischer TK. Rapid emergence and antigenic diversification of the norovirus 2012 Sydney variant in Denmark, October to December, 2012. *Euro Surveill.* 2013;18(9):pii=20413. Available online: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=20413>

⁴Bennett S, MacLean A, Miller RS, Aitken C, Gunson RN. Increased norovirus activity in Scotland in 2012 is associated with the emergence of a new norovirus GII.4 variant. *Euro Surveill.* 2013;18(2):pii=20349. Available online: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=20349>



The VIDRL Norovirus Report June 2013

VIDRL
Victorian Infectious Diseases
Reference Laboratory

Available at: www.vidrl.org.au

Figure 1: Norovirus detections by age group and year, VIDRL, January 2003 – June 2013




Figure 2: Numbers and rates of gastroenteritis consultations, MMDS, January 2003 – June 2013

